



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

of their environment. In short, it is a study of the ecological anatomy of fossil leaves. The leaves of *Calamites* are mostly known as impressions, and THOMAS has referred the structures he has been able to obtain to the better known impression forms. The petrified material used was obtained from the Lower Coal-measures (Halifax Hard Bed), and five types of leaf were distinguished. The twigs bearing the leaves proved to be quite interesting in stelar structure, especially in its relation to that of young stems of *Equisetum*.

The ecological conclusion is as follows: "The leafy twigs seem to have grown in a pendulous fashion, and the structure of the mesophyll and epidermis suggests that the habitat was a damp one. On the other hand, the leaves possess some xeromorphic features, such as the presence of fibers in the longer forms. The evidence points to a marsh or swamp forest as their habitat; this may have been near the sea, but if so the soil probably contained little salt."—J. M. C.

Transpiration in salt marsh plants.—Transpiration rates of cut shoots of *Salicornia annua* and *Suaeda maritima* have been found by DELF⁴³ to be equal to or greater than those from equal surface areas of such mesophytes as *Vicia Faba*. The highest degree of succulence seemed to be accompanied by the highest transpiration rate per unit area. Relating transpiration to evaporation from a water surface, unit areas of *Salicornia* lost 32 per cent and of *Vicia* 26 per cent as much water as equal areas of water surface. It was shown that *Salicornia* is able to absorb water readily through the surface of its stems when submerged, and less convincingly that it does not absorb sufficiently through its root system to replace the loss by transpiration except in a humid atmosphere. This seems the more surprising, since TRANSEAU⁴⁴ has shown evaporation to be exceptionally high in salt marshes. The stomata in *Salicornia* and *Aster trifolium* appear to have distinct powers of movement in young plants early in the season, losing this plasticity at a later date.—GEO. D. FULLER.

Cytology of the ascus.—An account of the cytology of *Helvella crispa* Fries is given by Miss CARRUTHERS.⁴⁵ The cells of the hypothecium are one to several-nucleate, and some of the nuclei were observed to fuse in pairs, but no migration was observed like that in *Humaria*. An attempt was made to determine the number of chromosomes on the spindles in the vegetative hyphae. Apparently two chromosomes are present in the vegetative spindles, and four or eight on the spindles in the fertile hyphae, but owing to the minuteness of the objects not much importance can be attached to these observations. The

⁴³ DELF, E. MARION, Transpiration and behavior of stomata in halophytes. Ann. Botany 25:484-505. 1911.

⁴⁴ BOT. GAZ. 45:217-231. 1908.

⁴⁵ CARRUTHERS, D., Contributions to the cytology of *Helvella crispa* Fries. Ann. Botany 25:243-252. pls. 2. 1911.